

PRODIAMINE GROUP 3 HERBICIDE



PROFESSIONAL TURF PRODUCTS

0.48% BARRICADE[®] ON DG PRO[®]



PRE-EMERGENT WEED CONTROL FOR LAWNS & LANDSCAPE ORNAMENTALS



Dispersing Granule Technology



Controls over 30 grassy and broadleaf weeds



Small particle formulation provides
more particles per square inch

ACTIVE INGREDIENT:
Prodiamine*.....0.48%
OTHER INGREDIENTS:.....99.52%
Total 100.00%
*(CAS No. 29091-21-2)

EPA Reg. No. 9198-214
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A,
9198-IN-3S
Underlined letter is first letter used in run code.

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE BACK PANEL AND BOOKLET FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND FIRST AID

NET WEIGHT 18.00 LBS (8.16 KG)
COVERS UP TO 5,800 SQ. FT.



PROFESSIONAL TURF PRODUCTS

0.48% BARRICADE® ON DG PRO®

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

WHERE TO USE

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens), lawn and sod nurseries
- container, field-grown, and landscape ornamentals
- established perennial and wildflower plantings
- christmas tree farms

HOW THIS PRODUCT WORKS

This product controls susceptible weeds by preventing growth and development of newly germinated weeds. Weed control is most effective when product is watered-in, within 2 weeks, by at least 0.5 inch of rainfall or irrigation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

Product of U.S.A.

MANUFACTURED BY:

The Andersons, Inc.
PO Box 119 | Maumee, OH 43537
info@andersonshomeandgarden.com
877-474-8262

QUESTIONS?

For additional product information, lawn care tips and spreader settings, visit:


AndersonsHomeandGarden.com

Find us on social media: @AndersonsHomeandGarden   

10007835
30001200
F22

APPLICATION DIRECTIONS**ESTABLISHED TURF/LAWNS**

1.) For best results this product must be applied prior to crabgrass germination. To determine the suitable timing for your specific region please consult the map below.



2.) Select a properly calibrated turf spreader and adjust the rate in accordance with the spreader settings listed below. Most established lawns can achieve season-long control of summer annual weeds at the low or medium rate listed below. For spreader models not located below, please refer to The Andersons website for further details.

SPREADER SETTINGS

Recommended Spreader Settings				
These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.				
18 lbs. treats 5,800 sq. ft. at 3.1 lbs/1,000 sq. ft. rate (Low Rate)				
18 lbs. treats 5,100 sq. ft. at 3.5 lbs/1,000 sq. ft. rate (Medium Rate)				
18 lbs. treats 3,800 sq. ft. at 4.7 lbs/1,000 sq. ft. rate (High Rate)				
18 lbs. treats 2,500 sq. ft. at 7.1 lbs/1,000 sq. ft. rate (Heavy Rate)				
Spreader	Spreader Settings			
	Low Rate	Medium Rate	High Rate	Heavy Rate
The Andersons Pacer Pro	8	10	12	14
The Andersons LCO-1000	K 1/2	L	M 1/2	P
The Andersons Yard Star Z150	13 1/2	14	15 1/2	18
Agri-Fab Broadcast	3 1/2	4	5	7
Brinly 20 Series Push Broadcast	13 1/2	14	15 1/2	18
Chapin Broadcast	13 1/2	14	15 1/2	18
Earthway Broadcast	13 1/2	14	15 1/2	18
Echo Broadcast	4	4 1/2	5 1/2	7 1/2
Lesco Broadcast	11 1/2	12	12 1/2	15
Scotts Broadcast*	4	4 1/2	5 1/2	7 1/2

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

3.) For best results the product should be activated with 0.5 inches of irrigation following application.

Do not apply this product to creeping bentgrass mowed to less than 0.5 inches in height.

FURTHER DETAILS ABOUT RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Depending on your turf type, please refer to the table below for the maximum amount of product that can be applied in a single growing season.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied. The low rate will provide 7 months of control and the medium rate will provide 8 months of control. Higher rates will control weed seeds longer but be careful as a longer residual will also hinder future over-seeding of desired turf species. Additionally, some species can be sensitive to dosage. Please refer to the maximum annual rates below.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.

MAXIMUM APPLICATION RATE OF THIS PRODUCT PER CALENDAR YEAR BY TURF GRASS SPECIES¹

Turf Species	Lbs. Ai/A	Lbs. Product/A	Lbs. Product/1,000 sq. ft.
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turftype), Zoysia	1.5	312	7.1
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	1.0	208	4.7
Creeping Red Fescue	0.75	156	3.5
Creeping Bentgrass	0.65	135	3.1

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 108 lb./acre or 2.4 lb./1,000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Use higher rates of this product to achieve higher levels weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

THE ANDERSONS PROFESSIONAL TURF PRODUCTS 0.48% BARRICADE ON DG PRO

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WPS USES:

Applicators and other handlers:

- Long sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash all PPE separately from other laundry.

NON-WPS USES:

Mixers and loaders who handles this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, only agricultural-plant uses are covered by the WPS – must wear:

- Chemical-resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash all PPE separately from other laundry.

ENGINEERING CONTROLS:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish.

Aquatic Organism Advisory Statement: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash waters.

If material is spilled, contain and sweep up material of spill for disposal or reuse per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under or cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

WEED RESISTANCE MANAGEMENT

For resistance management, this product is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading

patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance- management and/or integrated weed-management recommendations for specific crops and weed biotypes.

WEEDS CONTROLLED

When used at the recommended rates this product will control:

Barnyardgrass	Itchgrass	Pusley, Florida
Bluegrass, Annual ¹ (<i>Poa annua</i>)	Johnsongrass (from seed)	Rescuegrass ⁴
Carpetweed	Junglerice	Shepherd's Purse ³
Chickweed, Common	Knotweed	Signalgrass, Broadleaf
Chickweed, Mouseear (from seed)	Kochia	Speedwell, Persian
Crabgrass (Large, Smooth)	Lambsquarter, Common	Sprangletop
Crowfootgrass	Lovegrass	Spurge, Prostrate
Cupgrass, Woolly	Oxalis	Witchgrass
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Woodsorrel, Yellow (from seed)
Goosegrass ²	Pigweed	
Henbit	Purslane, Common	

¹ In areas where *Poa annua* is a winter annual, apply this product in approximately August or September to established, non-overseeded turf before *Poa annua* seeds germinate.

² In many areas a single application of 3.1 to 7.1 lbs./1,000 sq.ft. (135-312 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 3.1 to 4.7 lbs./1,000 sq. ft. (135-208 lbs./A) followed after 60 to 90 days by a application at doses that would not exceed those given in the Maximum Application Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Application Rates Table above.

³ Applications for this weed should be made in late summer, fall, or winter prior to germination.

⁴ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION - ALL STATES*

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

APPLICATION RATE FOR THIS PRODUCT			INTERVAL BEFORE OVERSEEDING		
Lbs./1,000 sq. ft.	Lbs./A	ai/A	North	Transition	South
2.3	104	0.50	4 mo.	4 mo.	4 mo.
3.1	135	0.66	5 mo.	4 mo.	4 mo.
3.5	156	0.74	6 mo.	5 mo.	5 mo.
3.8	166	0.80	-	6 mo.	6 mo.
4.7	208	1.00	-	7 mo.	7 mo.
5.4	237	1.14	-	-	9 mo.
6.1	270	1.30	-	-	10 mo.
7.1	312	1.50	-	-	12 mo.

* Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section "*Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV, and TX only)".

POA ANNUA CONTROL IN ESTABLISHED BERMUDAGRASS OVERSEEDED WITH PERENNIAL RYEGRASS (AZ, CA, NV, AND TX ONLY)

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

HOW MUCH AND WHEN TO APPLY

AMOUNT TO APPLY		WHEN TO APPLY	EXPECTED CONTROL	USE PRECAUTIONS
Lbs./Acre	Lbs./1,000			
78 - 135	1.8 - 3.1	6 to 8 weeks before ryegrass overseeding Second application 4-8 weeks after overseeding date or when perennial ryegrass roots are in the second inch of soil	One application for 70% or greater control of <i>Poa annua</i> Second application should enhance the residual under normal weather conditions	1. Some seed mortality and temporary reduction in root growth of new seedlings may occur to reduce the potential for seeding mortality, maintain a moist seed-bed with light, frequent irrigation. 2. Make no more than 2 applications per year for this use, and do not exceed 4 lbs./1,000 sq. ft. (176 lbs./A) for these two applications for <i>Poa annua</i> control in ryegrass overseeded turf 3. Do not make a second application if any injury to the ryegrass is observed after the first application 4. Do not make a second application unless the product was first applied before overseeding

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over seeded grasses.
- Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

USE RATES

Apply this product at 3.1 to 7.1 lbs./1,000 sq. ft. (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 sq. ft. (312 lbs./acres).

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Apply this product to dry foliage only. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate this product in the soil.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed on the next page are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Fir species** (Balsam, Fraser, Noble, etc.)
 Abelia: Sherwood
 Japanese Maple
 Norway Maple
 Yarrow: King Edward
 Kiwi*
 Lily-of-the-Nile (African Lily)
 Five-Leaf or Chocolate Vine
 Lady's Leek, Nodding Onion
 Japanese Anemone
 Aquilegia: Red and gold
 Vine Hill Manzanita
 Cape Weed
 Wormwood; Silver Mound, Castle
 Japanese Aucuba
 Barberr
 Aster: Bonny Blue, Purple Dome
 Lady Fern; Fern Lady
 Fibrous Begonia; Hardy Grandis
 Barberr
 Wintergreen Barberr
 Mentor Barberr
 Japanese Barberr
 Warty Barberr
 Snowbank
 Bougainvillea
 Butterfly-Bush (Dwarf Blue); Royal
 Red
 Japanese Boxwood
 Crimson Bottlebrush
 Weeping Bottlebrush
 Scotch Heather
 Tussock Bellflower; (White Clips)
 Trumpet Creeper, Trumpet Flower; Madame Galen
 Hottentot Fig (Ice Plant)
 Feathery Cassia
 Wild Lilac
 False Cypress
 Cleyera

Citrus species*
 Coreopsis (Calliopsis); Early Sunrise
 Moonbean
 Flowering Dogwood
 American Dogwood
 Pampas Grass
 Cranberry Cotoneaster
 Cotoneaster
 Bearberry Cotoneaster
 Rockspray Cotoneaster
 Hawthorn
 Lucifer
 Italian Cypress
 White Trailing Ice Plant
 Cooperi Pink
 Larkspur; Blue Elf
 Dianthus, Maiden Pinks 'Zing'
 Cheddar Pink
 Hop Bush
 Coneflower, Purple; Magnus
 Silverberry
 Wintercreeper
 Japanese Spindle Tree (Evergreen Euonymus)
 Spreading Euonymus
 Japanese Aralia
 Border Forsythia
 Weeping Forsythia
 Greenstem Forsythia
 Gallaria, Blanket Flower: 'Goblin'
 Gardenia, Cape-Jasmine; Radicans
 Siberian iris; Cabernet
 Iris species**
 Winter Jasmine
 Chinese Juniper; Nick's Compact,
 Parsoni
 Shore Juniper; Blue Pacific
 Parsoni
 Creeping Juniper
 Walnut*
 Shrimp Plant
 Grape Myrtle

Crape Myrtle; Tuscarora
 Weeping Lantana
 Lavender; Munstead
 Edleweiss
 Chinese Privet; Variegata
 Amur Privet
 Japanese Privet
 Glossy Privet (Wax-Leaf)
 Lily; Jazz
 Big Blue Lillyturf; Turf, Silver Mound,
 Evergreen Giant
 Liriope, Variegated
 Liriope, Creeping
 Cardinal Flower, Indian Pink
 Japanese Honeysuckle
 Tatarian Honeysuckle
 Burgundy
 Loosestribe; Modern Pink
 Magnolia species**
 Ice Plant
 Grabapple*
 Yaku Jima, Silberfeder
 Heavenly Bamboo
 Narcissus species**
 Oleander
 Evening Primrose
 Olive* Mondo Grass (Holly);
 Osmanthus (False Holly); Gulf Tide
 Trailing African Daisy
 Sourwood
 Tree Peony
 Fountain Grass (Dwarf)
 Avocado*
 Frasier's Photinia (Redtip)
 Dragonhead; False; Vivid
 Spruce species** (Colorado Blue, Norway, etc.)
 Lily-of-the-Valley Shrub
 Calabrian Pine
 Canary Island Pine
 Firethorn
 Bradford Pear spp.
 Oak, Shumard's Red
 Oak species

Indian Hawthorn
 Yedda Hawthorn
 Rhododendron (including Azalea)
 'Coral Bells'
 'Delaware Valley White'
 'Flame Creeper'
 'Formosa'
 'Girard Crimson'
 'George I. Tabor'
 'Hino-crimson'
 'PJM'
 'Roseum Elegans'
 Wakeibeisu
 White Gumbo
 Lady Bank's Rose
 Rosemary*
 Black-Eyed Susan; Goldstrum
 Leatherleaf Fern
 Santolina
 Saxifrage; Purple Dome
 Pincushion Flower
 Stonecrop; Lidakense, Dragon's
 Blood
 Spirea: Anthony Waterer
 Australian Brushcherry
 Japanese Yew
 Yew
 Germander
 Meadow Rue
 American Arborvitae
 Star Jasmine
 Canada Hemlock
 Tulip species
 Veronica Speedwell; Sunny
 Border
 Japanese Viburnum
 Sweet Viburnum
 Japanese Snowball
 Canary Island Viburnum
 Arrowwood Viburnum
 Laurustinus
 Cranberry Bush
 Leatherleaf Viburnum
 Greater Periwinkle, Vinca

Common Periwinkle, Vinca Grape*
 Old Fashioned Weigela
 Spanish Bayonet, Yucca
 Adam's Needle, Yucca
 Gentian
 Cranesbill
 Gladiolus species**
 Baby's Breath
 English Ivy
 Sunrose
 Daylily; Aztec Gold, Stella De Oro, Tender Love
 Coral Bell; Bridget Bloom
 Rose of Sharon**
 Chinese Hibiscus
 Mallow; Disco Belle White
 Hosta, Plantain Lily (Fragrant)
 Hosta, 'Searsucker'
 Bigleaf Hydrangea
 Chinese Holly
 Japanese Holly; Helleri
 American Holly
 Holly
 Yaupon Holly; Schillings
 Sword-Leaved Iris; Jodlesong
 Slash Pine
 Aleppo Pine
 Austrian Black Pine
 Longleaf Pine
 Monterey Pine
 Japanese Black Pine
 Eastern White Pine
 Scotch Pine
 Loblolly Pine
 Virginia Pine
 Pistachio*
 Queensland Pittosporum
 Japanese Pittosporum
 Japanese Yew
 English Laurel
 Almond, Apricot, Nectarine,
 Peach, Plum and Prune*
 Douglas Fir**
 Firethorn Scarlet

VEGETATION MANAGEMENT

This product may be applied to soil surfaces for preemergence control of many grass and broadleaf weeds in non-cropland areas on or surrounding:

- Non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways & equipment yards)
- Facilities including substations, tankfarms, pumping stations, parking and storage areas and ungrazed fencerows

Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Apply this product at 3.1 to 7.1 lbs./1,000 sq. ft. (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 sq. ft. (312 lbs./A).

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of U.S.A

Manufactured by:
 The Andersons, Inc. | P.O. Box 119 | Maumee, OH 43537

©2025 The Andersons, Inc. DG Pro is a registered trademark of The Andersons, Inc. Barricade is a registered trademark of Syngenta Crop Protection. **J25**

*Do not use on food producing trees, vines, or plants.

** Not for use on container grown plants.