

# Tee Time®

## SGN 100

# 0 + 0 + 30 + Fe, Mn, Zn, Cu

## (0 + 0 + 24.9 + Fe, Mn, Zn, Cu)

**USA/INT**      **0 + 0 + 30 + Fe, Mn, Zn, Cu**

30.0% Soluble Potash (K<sub>2</sub>O) from potassium sulfate  
 10.2% Total Sulfur (S) from sulfate  
 0.1% Copper (Cu) from copper oxide  
 3.0% Iron (Fe) from ferric oxide  
 0.1% Manganese (Mn) from manganous oxide  
 0.1% Zinc (Zn) from zinc oxide

**Application rates: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**

**Directions for use:** Apply product to actively growing turf at 15–25 g/m<sup>2</sup>. Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.

**UK/EC**      **EC FERTILISER**  
**POTASSIC FERTILISER WITH Fe, Mn, Zn, Cu**  
**0 + 0 + 30 + Fe, Mn, Zn, Cu**

30.0% Potassium oxide (K<sub>2</sub>O) water soluble  
 25.5% Sulphur Trioxide (SO<sub>3</sub>) water soluble  
 0.1% Copper (Cu)  
 3.0% Iron (Fe)  
 0.1% Manganese (Mn)  
 0.1% Zinc (Zn)

Low in chloride. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.  
 Import in U.K. by: Greentech Sportsturf Ltd.  
 Block 5 Unit 1 - Banded Industrial Estate  
 Throsk, Stirling - Scotland FK7 7NP

**Application rates: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**  
**Directions for use:** Apply product to actively growing turf at 15–25 g/m<sup>2</sup>. Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.

**Eire/EC**      **EC FERTILISER**  
**POTASSIC FERTILISER WITH Fe, Mn, Zn, Cu**  
**0 + 0 + 24.9 + Fe, Mn, Zn, Cu**

24.9% Potassium (K) water soluble  
 10.2% Sulphur (S) water soluble  
 0.1% Copper (Cu)  
 3.0% Iron (Fe)  
 0.1% Manganese (Mn)  
 0.1% Zinc (Zn)

Low in chloride. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.  
 Import in EIRE by: Greentech Sportsturf Ltd.  
 Block 5 Unit 1 - Banded Industrial Estate  
 Throsk, Stirling, Scotland FK7 7NP

**Application rates: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**  
**Directions for use:** Apply product to actively growing turf at 15–25 g/m<sup>2</sup>. Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.

**New Zealand**      **0 + 0 + 24.9 + Fe, Mn, Zn, Cu**

24.9% Potassium (K) from potassium sulfate  
 10.2% Total Sulfur (S) from sulfate  
 0.1% Copper (Cu) from copper oxide  
 3.0% Iron (Fe) from ferric oxide  
 0.1% Manganese (Mn) from manganese oxide  
 0.1% Zinc (Zn) from zinc oxide

**Application rates: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**

**Directions for use:** Apply product to actively growing turf at 15–25 g/m<sup>2</sup>. Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.

**Deutschland**  
**K-Dünger 30, mit Schwefel und Spurennährstoffen zur Düngung von Rasen Für Kinder unzugänglich aufbewahren!**

30,0 % K<sub>2</sub>O wasserlösliches Kaliumoxid  
 10,2 % S wasserlöslicher Schwefel  
 0,1 % Cu Gesamt-Kupfer  
 3,0 % Fe Gesamt-Eisen  
 0,1 % Mn Gesamt-Mangan  
 0,1 % Zn Gesamt-Zink

Mit gebrochenem Kalkstein als Ausbringhilfe.  
 Nettogewicht: 20 kg

**Inverkehrbringer:** Turf Handels GmbH, Am Hartboden 48, A-8101 Gratkorn

**Aufwandmenge: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**

**Anwendungsempfehlung:** Während der Vegetationszeit mit 15–25 g/m<sup>2</sup> auf trockene Bestände ausbringen. Anschließend mindestens mit 5–6 mm beregnen, damit die Körner gleichmäßig in die Rasenarbe fallen. Bei Temperaturen über 25°C nur mit geringer Aufwandmenge streuen und sofort beregnen. Das Produkt enthält Eisen, das auf gepflasterten Oberflächen im Zusammenhang mit Wasser Flecken verursachen kann. Sollte Dünger auf derartige Oberflächen gelangen, nicht mit Wasser entfernen, sondern sofort abkehren. Trocken und sauber lagern, Restmengen dicht und trocken verschließen. Nur saubere Streuer benutzen, keine Restmengen im Streuwagen zurücklassen. Um Aufnahme des Düngers zu vermeiden, den ersten Schnitt ohne Fangkörbe durchführen.

**Österreich**  
**Mineralischer Kalidünger, mit Spurennährstoffen 0-0-30 Für Kinder unzugänglich aufbewahren!**

30,0 % K<sub>2</sub>O wasserlösliches Kaliumoxid  
 10,2 % S wasserlöslicher Schwefel  
 0,1 % Cu Gesamt-Kupfer  
 3,0 % Fe Gesamt-Eisen  
 0,1 % Mn Gesamt-Mangan  
 0,1 % Zn Gesamt-Zink

Komponenten: Kaliumsulfat, Kupferoxid, Kupfersulfat, Eisenoxid, Eisensulfat, Manganoxid, Mangansulfat, Zinkoxid, Zinksulfat, Kalkstein.

Nettogewicht: 20 kg  
 Hersteller: Turf Handels GmbH, Am Hartboden 48, A-8101 Gratkorn

**Aufwandmenge: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**  
**Anwendungsempfehlung:** Während der Vegetationszeit mit 15–25 g/m<sup>2</sup> auf trockene Bestände ausbringen. Anschließend mindestens mit 5–6 mm beregnen, damit die Körner gleichmäßig in die Rasenarbe fallen. Bei Temperaturen über 25°C nur mit geringer Aufwandmenge streuen und sofort beregnen. Das Produkt enthält Eisen, das mit Wasser Flecken auf Pflastern und Platten hinterlassen kann. Wenn Düngerpartikel auf andere Oberflächen gelangen, unmittelbar nach dem Streuen durch Kehren oder Absaugen entfernen. Auf keinen Fall Abregnen, da es im Kombination mit Wasser zu Flecken kommt. Trocken und sauber lagern, Restmengen dicht und trocken verschließen. Nur saubere Streuer benutzen, keine Restmengen im Streuwagen zurücklassen. Um Aufnahme des Düngers zu vermeiden, den ersten Schnitt ohne Fangkörbe durchführen.

**Italia**      **CONCIME CE**  
**0 + 0 + 30 + Fe, Mn, Zn, Cu**

30,0% Ossido di potassio (K<sub>2</sub>O) solubile in acqua  
 0,1% Rame (Cu)  
 3,0% Ferro (Fe)  
 0,1% Manganese (Mn)  
 0,1% Zinco (Zn)  
 25,5% Anidride solforica (SO<sub>2</sub>) solubile in acqua

Basso contenuto di cloro. Da usare solo quando vi è un bisogno riconosciuto. Non superare i dosaggi appropriati.  
 Prodotto e confezionato da: The Andersons, Inc., Maumee - 43537, OH (USA)  
 Importato (fabbricante) da: BOTTOS s.r.l.  
 via Lusevera,1 – 33078 San Vito al Tagliamento (PN)

Distribuito da: EMERALD s.r.l.  
**Dose:** 15–25 g/m<sup>2</sup> (150–250 kg/ha)  
**Indicazioni per l'uso:** applicare 15–25 g/m<sup>2</sup> di prodotto su manto erboso in crescita. Applicare su manto asciutto. Irrigare dopo l'applicazione coprendo il manto erboso con 5 mm o 6 mm di acqua. Se la temperatura è superiore ai 25°C o se si è forzati ad applicare il prodotto su manti umidi o bagnati, irrigare immediatamente dopo l'applicazione. Applicare il prodotto a basse concentrazioni se la temperatura eccede i 25°C. Il presente prodotto contiene ferro, che potrebbe macchiare superficiali in cemento o altri materiali. Per evitare che il prodotto macchi, rimuovere le particelle immediatamente o usando una scopa o un soffiatore. NON PULIRE USANDO ACQUA. Conservare in luogo asciutto e pulito. Risigillare il sacchetto aperto piegando ed assicurando la parte superiore. Usare solo spargitori puliti ed asciutti. Non lasciare prodotto non usato negli spargitori.

**Norge**      **K-gjødsel med Fe, Mn, Zn, Cu**  
**0 – 0 – 25**  
**(0 – 0 – 30)**

24,9% Kalium (K)  
 30,0% kaliumoksyd (K<sub>2</sub>O)  
 10,2% Svovel (S)  
 3,0% Jern (Fe)  
 0,1% Kobber (Cu)  
 0,1% Mangan (Mn)  
 0,1% Sink (Zn)

Basert på kalium-, jern-, kobber-, mangan- og sink-sulfat, samt jern-, kobber-, mangan- og sinkoksyd. Skal bare benyttes ved påvist behov til golf- og fotballbaner, park- og hagearealer. De anbefalte mengder må ikke overskrides.  
 Distributør i Norge: Floratine Norge AS  
 Gullfunnet 50  
 1570 Dilling, NORGE

**Anbefalte gjødselmengder: 15–25 g/m<sup>2</sup> (150–250 kg/ha)**

Gjødselia tilføres når grasen er i vekst. Det vannes 4-5 minutter etter at gjødsling er utført, for å få gjødselkornene ned i plantebestandet. Bruk mindre gjødselmengder ved høge temperaturer (> 25°C), og vann øyeblikkelig for å unngå sviskader. Dette produktet inneholder jern som kan misfarge betong og andre overflater. Fjern partikler omgående ved feing eller blåsning. VASK IKKE BORT MED VANN. I forbindelse med gjødsling skal det kun benyttes tørre og rene gjødselspredere. La ikke gjødsel bli liggende igjen i sprederen. Gjødselia skal lagres tørt.

**Canada**      **0 + 0 + 30 + Fe, Mn, Zn, Cu**  
**GUARANTEED MINIMUM ANALYSIS**

30.0% Soluble Potash (K<sub>2</sub>O)  
 10.2% Sulfur (S)  
 7.0% Calcium (Ca)  
 0.1% Copper (Cu)(actual)  
 0.02% Water Soluble Copper (Cu)  
 3.0% Iron (Fe)(actual)  
 0.03% Water Soluble Iron (Fe)  
 0.1% Manganese (Mg)(actual)  
 0.03% Water Soluble Manganese (Mn)  
 0.1% Zinc (Zn)(actual)  
 0.03% Water Soluble Zinc (Zn)

Registration Number XXXXXXXM ANALYSE MINIMALE GARANTIE  
 Fertilizers Act  
 Lot# xxxxxxxx

**Application rates: 1.23 - 2.03 kg/100 m<sup>2</sup> (2.50 - 4.15 lbs/1,000 sq. ft.)**  
**Directions for use:** Seek advice from the county agricultural representative or a professional agricultural consultant. This product may not provide all nutrients required and should be used as part of a complete fertilization program, based on soil testing.  
**Caution:** this fertilizer contains copper, iron, manganese and zinc and should be used only as recommended. It may prove harmful when misused.

**WARNING**  
 MAY CAUSE EYE IRRITATION  
 MAY CAUSE RESPIRATORY IRRITATION

**PERSONAL PROTECTIVE EQUIPMENT / PROTECTION MEASURES / CONTROLS:** RESPIRATORY PROTECTION: NIOSH approved particulate respirator, if required. EYE PROTECTION: Safety glasses with side shields, goggles, or face shield recommended. SKIN PROTECTION: Long sleeves, chemical-resistant gloves recommended. VENTILATION: Local exhaust ventilation recommended.  
**PRECAUTIONARY STATEMENTS:** Call a poison center / physician if unwell. Wash hands after handling. Use only outdoors or in well-ventilated areas. Avoid breathing dust. Store container tightly closed. Keep out of reach of children. Dispose of contents / container in accordance with national / regional / local regulations.  
**FIRST AID:** IF INHALED: Move victim to fresh air. Seek medical attention if irritation persists. IF ON SKIN: Wash affected areas with soap and water. Seek medical attention if irritation persists. Wash contaminated clothing before re-use. IF IN THE EYES: Immediately flush with water for at least 20 minutes. Seek medical attention if irritation persists. IF SWALLOWED: If victim is alert and not convulsing, give one glass of water to dilute material but DO NOT induce vomiting. Seek immediate medical attention.  
**PREMIERS SOINS:** EN CAS D'INHALATION: Déplacez la victime à l'air frais. Consulter un médecin si l'irritation persiste. SI SUR LA PEAU: Lavez les zones touchées avec de l'eau et du savon. Consulter un médecin si l'irritation persiste. Laver les vêtements contaminés avant de les réutiliser. SI DANS LES YEUX: Rincer immédiatement à l'eau pendant au moins 20 minutes. Consulter un médecin si l'irritation persiste. EN CAS D'INGESTION : Si la victime est alerte et ne convulse pas, donnez un verre d'eau pour diluer le matériel, mais NE PAS provoquer de vomissements. Consulter immédiatement un médecin.

**Taiwan (R.O.C.)**      **菁好牌鉀霸 (0-0-30)**

肥料登記證字號: 肥進(鉀)字第 0128026 號  
 肥料品目: 粗製鉀鹽肥料(3-0-4)  
 國外廠牌商品名稱: TURF FERTILIZER 0-0-30  
 保證成份: 水溶性氧化鉀30.0%  
 成品性狀: 固態 顆粒狀  
 使用方法: 一般用法為春天至秋末使用 六至八週使用一次 施用後澆水10-20分鐘  
 使用量: 每平方公尺20公克 (每分鐘20-30 kg)  
 包裝方式: 袋裝      淨重: 20 kg  
 有效期限: 4年      批號: 請詳見包裝  
 製造日期: 請詳見包裝      電話: 04-22135266  
 總代理: 菁修企業有限公司  
 地址: 台中市東區自由路四段210號  
 製造地: 美國  
 製造廠商: THE ANDERSONS, INC.  
 P. O. BOX 119, MAUMEE, OH USA 43537

**Taux d'application: 1,23 – 2,03 kg/100 m<sup>2</sup> (2,50 – 4,15 lb/1 000 pP)**  
**MODE D'EMPLOI:** Demandez conseil au représentant agricole du comté ou à un consultant agricole professionnel. Ce produit peut ne pas fournir tous les éléments nutritifs requis et doit être utilisé dans le cadre d'un programme complet de fertilisation, basé sur des analyses de sol.  
**Attention:** cet engrais contient du cuivre, du fer, du manganèse et du zinc et ne doit être utilisé que selon les recommandations. Il peut s'avérer nocif lorsqu'il est mal utilisé.

**AVERTISSEMENT**  
 PEUT CAUSER UNE IRRITATION DES YEUX  
 PEUT CAUSER UNE IRRITATION RESPIRATOIRE

**ÉQUIPEMENT DE PROTECTION INDIVIDUELLE / MESURES DE PROTECTION / CONTRÔLES:** PROTECTION RESPIRATOIRE: Respirateur à particules approuvé par le NIOSH, au besoin. PROTECTION DES YEUX: Lunettes de sécurité avec écrans latéraux, lunettes de protection ou écran facial recommandé. PROTECTION DE LA PEAU: Manches longues, gants résistant aux produits chimiques recommandés. VENTILATION: Ventilation par aspiration locale recommandée. **MISES EN GARDE:** Appelez un centre antipoison / médecin si vous ne êtes pas bien. Se laver les mains après la manipulation. Utiliser uniquement à l'extérieur ou dans des endroits bien ventilés. Evitez de respirer de la poussière. Entroposer le contenant hermétiquement fermé. Tenir hors de la portée des enfants. Éliminer le contenu / le conteneur conformément aux réglementations nationales / régionales / locales.

**Australia**      **0 + 0 + 24.9 + Fe, Mn, Zn, Cu**

24.9% Potassium (K) from potassium sulfate  
 10.2% Total Sulfur (S) from sulfate  
 0.1% Copper (Cu) from copper oxide  
 3.0% Iron (Fe) from ferric oxide  
 0.1% Manganese (Mn) from manganese oxide  
 0.1% Zinc (Zn) from zinc oxide

**Application rates: 1.5–2.5 kg/100 m<sup>2</sup> (150–250 kg/ha)**

**Directions for use:** Apply product to actively growing turf at 1.5–2.5 kg/100 m<sup>2</sup>. Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.

Spreaders	Ground Speed	Width of Coverage	Spreader Settings		
			15 g/m <sup>2</sup>	20 g/m <sup>2</sup>	25 g/m <sup>2</sup>
Model 2000 (cone 9); 2000 SR (cone 9) Settings for single pass application = FULL RATE	5 km/hr	3.7 m	1 3/4	J 3/4	L
SSD Drop Spreader Settings for single pass application = FULL RATE	5 km/hr	0.9 m	3 1/2	4 1/4	5
Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.					



The Andersons logo and Tee Time are registered trademark of The Andersons, Inc. J25

Manufactured by (Fabriqué par):  
 The Andersons, Inc.  
 P. O. Box 119  
 Maumee, OH 43537  
 U.S.A.  
 AndersonsPro.com

**20 kg/net**  
**10009131**