

# PolySports™

## SGN 150

# 10 + 18 + 10

## (10 + 7.8 + 8.3)

# 30% NS-54®

## With HumicDG™

USA/INT	10 + 18 + 10
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen*
18.0%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
10.0% 3.69%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate Sulfur (S) 0.29% Free Sulfur (S) 3.40% Combined Sulfur (S)
*3.00% Slowly available nitrogen from polymer coated sulfur coated urea.	
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

UK/EC	EC FERTILISER NPK FERTILISER 10 + 18 + 10
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen
18.0%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
10.0%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in U.K. by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Bandeath Industrial Estate Throsk, Stirling Scotland FK7 7NP
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Eire/EC	EC FERTILISER NPK FERTILISER 10 + 7.8 + 8.3
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen
7.8%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
8.3%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in EIRE by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Bandeath Industrial Estate Throsk, Stirling Scotland FK7 7NP
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Italia	FERTILIZZANTE EC FERTILIZZANTE NPK 10 + 18 + 10
10,0%	Totale azoto (N) 3,81% Azoto ammoniacato 6,19% Azoto ureico
18,0%	Fosfato disponibile (P <sub>2</sub> O <sub>5</sub> ) dal fosfato di ammonio
10,0%	Potassa solubile (K <sub>2</sub> O) dal solfato di potassio
CONTIENE INGREDIENTI NON FERTILIZZANTI VEGETALI Acido umico (derivato da leonardite)..... 16%	
Prodotto e confezionato da: The Andersons, Inc., Maumee - 43537, OH (USA)	
Importato (fabbricante) da: BOTTOS s.r.l. via Lusevera, 1 - 33078 San Vito al Tagliamento (PN)	
Distribuito da: EMERALD s.r.l.	
Tassi di applicazione: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Istruzioni per l'uso: 20-30 g/m <sup>2</sup> di prodotto su manto erboso in crescita. Applicare su manto asciutto. Irrigare dopo l'applicazione coprendo il manto erboso con 5 mm o 6 mm di acqua. Se la temperatura eccede i 25°C applicare solo dosaggi bassi ed irrigare immediatamente dopo l'applicazione. Applicare il prodotto a basse concentrazioni se la temperatura eccede i 25°C. Non usare su putting greens in campi da golf. Conservare in luogo asciutto e pulito. Risigillare il sacchetto aperto piegando ed assicurando la parte superiore. Usare solo spargitori puliti ed asciutti. Non lasciare prodotto non usato negli spargitori.	

Australia	10 + 18 + 10
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen
18.0%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
10.0%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in U.K. by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Bandeath Industrial Estate Throsk, Stirling Scotland FK7 7NP
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Taiwan (R.O.C.)	EC FERTILISER NPK FERTILISER 10 + 18 + 10
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen
18.0%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
10.0%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in U.K. by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Bandeath Industrial Estate Throsk, Stirling Scotland FK7 7NP
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

New Zealand	10 + 18 + 10
10.0%	Total Nitrogen (N) 3.81% Ammoniacal Nitrogen 6.19% Urea Nitrogen
18.0%	Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) from ammonium phosphate
10.0%	Soluble Potash (K <sub>2</sub> O) from potassium sulfate
CONTAINS NON-PLANT FOOD INGREDIENT Humic Acid (derived from leonardite)..... 16%	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in U.K. by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Bandeath Industrial Estate Throsk, Stirling Scotland FK7 7NP
Application rates: 20-30 g/m <sup>2</sup> (200-300 kg/ha)	
Directions for use: Apply product to actively growing turf at 20-30 g/m <sup>2</sup> . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. Do not use on putting greens. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

--

--

--

--

--

--

--

--

--

 Spreaders	Ground Speed	Width of Coverage	Spreader Settings		
			20 g/m <sup>2</sup>	25 g/m <sup>2</sup>	30 g/m <sup>2</sup>
Model 2000 & 2000 SR (cone 6) Settings for single pass application = FULL RATE	5 km/hr	3.0 m	1 3/4	N	0
Lely Models WTR, WFR, HR & 1250 (pattern III-C) Settings for single pass application = FULL RATE	7 km/hr	9.7 m	5 1/2	6	6 1/2
Vicon (03 Series) Settings for single pass application = FULL RATE	7 km/hr	8.5 m	24	29	33
Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.					



Manufactured by:  
The Andersons, Inc.  
P. O. Box 119  
Maumee, OH 43537  
U.S.A.  
AndersonsPro.com

# 20 kg/net

# 10005445

J25 

The Andersons logo and NS-54 are registered trademarks of The Andersons, Inc.; PolySports and Humic DG are trademarks of The Andersons, Inc.