

# 25-3-10

## Fertilizer with NS-54<sup>®</sup> and DG Iron Sucrate<sup>™</sup>

- Contains NS-54<sup>®</sup> for extended feeding.
- Contains iron to help prevent deficiencies and heighten initial color response.

### Guaranteed Analysis

Total Nitrogen (N).....	25.00%
0.60% Ammoniacal Nitrogen	
24.40% Urea Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	3.00%
Soluble Potash (K <sub>2</sub> O).....	10.00%
Sulfur (S).....	5.61%
2.21% Free Sulfur (S)	
3.40% Combined Sulfur (S)	
Iron (Fe).....	2.00%

Derived from ammonium phosphate, iron sucrate, polymer coated sulfur coated urea and potassium sulfate.

\*23.82% slowly available urea nitrogen from polymer coated sulfur coated urea.

F002091

### Florida Applications:

We recommend that you follow the Florida Green Industries BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04706>

We recommend that you follow the Florida Golf Course BMP's at:

<http://www.flrules.org/Gateway/reference.asp?No=Ref-04702>

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

**Note:** This product contains the micronutrient iron for extra greening and to help prevent iron deficiency. Iron may stain concrete and other surfaces. Misapplied product to these surfaces should be removed promptly by sweeping or blowing. DO NOT WASH OFF WITH WATER.



**WARNING**  
CAUSES SKIN, RESPIRATORY AND SERIOUS EYE IRRITATION

### PRECAUTIONARY STATEMENTS:

Wear protective gloves and clothing. Remove contaminated clothing and wash before reuse. Do not breathe dusts. Use only outdoors or in well-ventilated area. Call poison control or seek medical attention if you feel unwell. Wash thoroughly after handling. Do not eat, drink or smoke when using this product.

### FIRST AID

If inhaled: Move to fresh air and keep comfortable, seek medical attention if unwell.

If on skin: Wash affected areas with soap and water. Seek medical attention if irritation persists. Wash contaminated clothing before re-use.

If in the eyes: Immediately flush with water for 15-20 minutes, remove contact lenses if present and easy to do – continue rinsing. Seek medical attention if irritation persists.

If swallowed: Rinse mouth. DO NOT INDUCE VOMITING unless directed by a medical professional. Seek medical attention if unwell.

### DIRECTIONS FOR USE

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

**Nitrogen Notice:** For those states with nitrogen application restrictions, including but not limited to MD, NH, NJ, PA and VA, do not apply more than 0.9 lbs. of nitrogen per 1,000 sq.ft.

**Phosphate Notice:** This lawn/turf fertilizer contains phosphorus and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorus deficiency. This fertilizer is not for routine maintenance of turf.

In Maryland, this product is providing nutrients in accordance with a soil test that was conducted by a laboratory identified under 8-803.7 of this article and performed no more than 3 years before the application.

### Application Rates:

LIGHT:	3.0 lbs/1,000 sq. ft.
MEDIUM:	4.0 lbs/1,000 sq. ft.
HIGH:	5.0 lbs/1,000 sq. ft.

### For all applications:

- Apply to dry foliage.
- Water in after application with a minimum of 0.2 inches to help move particles into the turf canopy.
- Do not collect clippings for at least one mowing following application to avoid particle pick-up.
- Do not use on putting greens.
- Do not use with a drop-type spreader.

### Cool Season Grasses:

- Apply at the MEDIUM or HIGH RATE in the spring and fall months.
- Apply at the LIGHT RATE in summer months or when temperatures exceed 85°F.

### Warm Season Grasses:

- Apply at the HIGH RATE during the growing season.
- Apply at the MEDIUM RATE when temperatures exceed 90°.

### Storage and Use

Store in a clean, dry place.  
Reseal opened bag by folding top down and securing.  
Use only in clean, dry spreaders.  
Do not leave unused product in spreaders.

### IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

### WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS.

### LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of U.S.A.

Manufactured by:  
The Andersons, Inc.  
P.O. Box 119  
Maumee, OH 43537  
AndersonsPro.com

The Andersons logo and NS-54 are registered trademarks of The Andersons, Inc. DG Iron Sucrate is a trademark of The Andersons, Inc.

SUGGESTED SPREADER SETTINGS					
Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader prior to applying product.					
50 lbs. feeds 16,600 sq. ft. (0.38 acre) with 0.75 lbs. N/1,000 sq. ft. at the LIGHT RATE					
50 lbs. feeds 12,500 sq. ft. (0.29 acre) with 1.00 lbs. N/1,000 sq. ft. at the MEDIUM RATE					
50 lbs. feeds 10,000 sq. ft. (0.23 acre) with 1.25 lbs. N/1,000 sq. ft. at the HIGH RATE					
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			LIGHT RATE 3.0 lbs/1,000 sq. ft.	MEDIUM RATE 4.0 lbs/1,000 sq. ft.	HIGH RATE 5.0 lbs/1,000 sq. ft.
Andersons 2000 series (cone 4); Andersons LCO-1000	3 mph	11 feet	L	M 1/4	N 1/2
Scotts R-8, R-8A (cone 4)	3 mph	11 feet	K	L 1/4	M 1/2
Lesco 020093 (pattern 0.25)	3 mph	8 feet	E	F	F 1/2
Lesco 705698 (pattern 0.25)	3 mph	8 feet	12	14	15
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern II-C	4.5 mph	40 feet	5 1/2	6 1/4	7
Vicon (03 Series)	4.5 mph	32 feet	25	30	34

**SGN Standard**  
**Net Wt. 50 Lbs. (22.68 kg)**



**10009473**