

Tee Time®

SGN 100

8 + 0 + 16 + Mg, Fe, Mn

(8 + 0 + 13.3 + Mg, Fe, Mn)

USA/INT	8 + 0 + 16 + Mg, Fe, Mn
8.0%	Total Nitrogen (N)
	8.0% Ammoniacal Nitrogen
16.0%	Soluble Potash (K ₂ O) from potassium sulfate
3.0%	Magnesium (Mg) from magnesium oxide
14.5%	Total Sulfur (S) from sulfate
2.0%	Iron (Fe) from iron oxide
5.0%	Manganese (Mn) from manganese oxide
Application rates: 15–30 g/m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 15-30 g/m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

UK/EC	EC FERTILISER NK FERTILISER CONTAINING Mg, Fe, Mn 8 + 0 + 16 + Mg, Fe, Mn
8.0%	Total nitrogen (N)
	8.0% Ammoniacal nitrogen (N)
16.0%	Potassium oxide (K ₂ O) water soluble
5.0%	Magnesium oxide (MgO)
36.25%	Sulphur trioxide (SO ₃) water soluble
2.0%	Iron (Fe)
5.0%	Manganese (Mn)
Low in chloride. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in U.K. by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Banded Industrial Estate Throsk, Stirling - Scotland FK7 7NP
Application Rates: 15–30 g/m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 15–30 g/m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Eire/EC	EC FERTILISER NK FERTILISER CONTAINING Mg, Fe, Mn 8 + 0 + 13.3 + Mg, Fe, Mn
8.0%	Total nitrogen (N)
	8.0% Ammoniacal nitrogen (N)
13.3%	Potassium (K) water soluble
3.0%	Magnesium (Mg)
14.5%	Sulphur (S) water soluble
2.0%	Iron (Fe)
5.0%	Manganese (Mn)
Low in chloride. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	
Import in EIRE by:	Greentech Sportsturf Ltd. Block 5 Unit 1 - Banded Industrial Estate Throsk, Stirling - Scotland FK7 7NP
Application rates: 15–30 g/m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 15–30 g/m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

France/EC	ENGRAIS EC ENGRAIS NK AVEC Mg, Fe, Mn 8 + 0 + 16 + Mg, Fe, Mn
8,0%	Azote totale (N)
	8,0% Azote ammoniacal (N)
16,0%	Oxyde de potassium (K ₂ O) soluble dans l'eau
5,0%	Oxyde de magnésium (MgO)
36,25%	Trioxyde de soufre (SO ₃)
2,0%	Fer (Fe)
5,0%	Manganèse (Mn)
Faible teneur en chlorure. À utiliser uniquement en cas de besoin reconnu. Ne pas excéder les taux de dosage appropriés.	
Importé en France par:	LOB GREEN S.A.S. 25 Rue de la Chapelle GOURNAY, 62560 VERCHOCQ - France
Taux d'application : 15–30 g/m² (150–300 kg/ha)	
Mode d'emploi : Appliquer le produit sur le gazon en plaques en état de croissance active à un taux de 15-30 g/m ² . Appliquer sur le feuillage sec. Arroser après application avec 5 à 6 mm d'irrigation. Dans des températures supérieures à 25°C ou si l'application pour humidifier ou mouiller le gazon est inévitable, irriguer immédiatement après application. Appliquer à de faibles taux si les températures sont supérieures à 25°C. Ce produit contient du fer, susceptible de tacher le béton et d'autres surfaces. Éliminer rapidement les particules en balayant ou en soufflant pour éviter toute tache. NE PAS LAYER À L'EAU. Entreposer dans un endroit propre et sec. Refermer hermétiquement le sac ouvert en repliant le haut et en sécurisant. Utiliser uniquement des épanduses propres et sèches. Ne pas laisser de produit non utilisé dans les épanduses.	

Deutschland/Österreich	EC-DÜNGER NK-DÜNGER MIT MAGNESIUM, EISEN UND MANGAN 8 + 0 + 16 + Mg, Fe, Mn
8,0 %	N Gesamtstickstoff
	8,0% N Ammoniumstickstoff
16,0 %	K ₂ O wasserlösliches Kaliumoxid
3,0%	Mg Magnesium
14,5%	S Schwefel
2,0 %	Fe Eisen
5,0%	Mn Mangan
Nettogewicht: 20kg	
Hersteller: Turf Handels GmbH, Am Hartboden 48, A-8101 Gratkorn	
Aufwandmenge: 15–30 g/m² (150–300 kg/ha)	
Anwendungsempfehlung: Das Produkt mit einer Streueinstellung von 15-30 g/m ² auf einem aktiv wachsenden, trockenen Rasen ausstreuen. Nach dem Ausstreuen mit 5 bis 6 mm Wasser bewässern. Bei Temperaturen von über 25 °C nur in geringen Mengen auftragen und sofort nach dem Ausstreuen bewässern. Dieses Produkt enthält Eisen, das Flecken auf Zement und anderen Oberflächen hinterlassen kann. Restpartikel sofort durch Wegfegen oder -blasen entfernen, um Fleckenbildung zu verhindern. NICHT MIT WASSER ABWASCHEN. An einem sauberen, trockenen Ort lagern. Den geöffneten Beutel durch Umklappen der Oberkante und Sichern verschließen. Nur saubere, trockene Streuer verwenden. Kein unverbrauchtes Produkt im Streuer lassen. Für Kinder unzugänglich aufbewahren!	

España/CE	ABONO CE NK abono que contenga Mg, Fe, Mn 8 + 0 + 16 + Mg, Fe, Mn
8,0%	Nitrógeno total (N)
	8,0% Nitrógeno amoniacal (N)
16,0%	de óxido de potasio (K ₂ O) soluble en agua
5,0%	de óxido de magnesio (MgO)
36,25%	de azufre (SO ₃) soluble en agua
2,0%	Hierro (Fe)
5,0%	Manganeso (Mn)
Bajo en cloruro. Para ser utilizado sólo cuando hay una necesidad reconocida. No sobrepasar las dosis recomendadas.	
Importador para España: Dosis de aplicación: 15–30 g/m² (150–300 kg/ha)	
Instrucciones de uso: Aplicación de productos en crecimiento activo del césped. Aplicar cuando la hoja esté seca. Regar después de la aplicación con 5 a 6 mm de agua de riego. Se aconseja aplicar sólo a dosis bajas si las temperatura es superior a 25°C. Regar inmediatamente después de la aplicación. Este producto contiene hierro que puede manchar zonas aplicadas u otras superficies. Eliminar las partículas sobrantes barriendo o soplando (máquinas sopladoras) para evitar manchas. NO SE LAVA CON AGUA. Conservar el envase en un lugar limpio y seco. Una vez abierto el saco evitar que entre humedad, asegurarse de cerrar la parte superior del saco. Utilice sólo abonadoras limpias y secas. No deje de producto no utilizado en las abonadoras.	

Italia	CONCIME CE Concime NK (S) 8–16 (36,2) contenente Azoto ammoniacale a basso tenore di cloro ottenuto per miscelazione
Azoto (N) totale	8 %
di cui Azoto (N) ammoniacale	8 %
Ossido di potasio (K ₂ O) solubile in acqua	16 %
Anidride solforica (SO ₃) solubile in acqua	36,25 %
Magnesio (MgO)	5 %
Ferro (Fe)	2 %
Manganese (Mn)	5 %
Importato (fabbricante) da: BOTTOS s.r.l. via Lusevera,1 – 33078 San Vito al Tagliamento (PN)	
Distribuito da: EMERALD s.r.l.	
Dose: 15–30 g/m² (150–300 kg/ha)	
Istruzioni per l'uso: Applicare 15-30 gr/mq di prodotto su manto erboso asciutto in crescita. Irrigare dopo l'applicazione con 5-6 mm di acqua. Se la temperatura eccede i 25 °C applicare a dosaggi bassi e irrigare immediatamente. Il presente prodotto contiene ferro, che potrebbe macchiare superfici in cemento o altri materiali. Per evitare che il prodotto macchi, rimuovere le particelle immediatamente o usando una scopa o un soffiatore. NON PULIRE USANDO ACQUA. Conservare in luogo asciutto e pulito. Risigillare il sacchetto aperto piegando ed asciugando la parte superiore. Usare solo spargitori puliti e asciutti. Non lasciare prodotto inusato negli spargitori.	

Sverige	EU GÖDSELMEDEL NK-gödselmedel med Mg och Mn 8 – 0 – 13 (8 – 0 – 16)
8,0 %	Kväve (N)
	8,0% ammonium-N
13,3 %	Kalium (K)
	16,0% kaliumoxid (K ₂ O)
3,0%	Magnesium (Mg)
14,5%	Svavel (S)
2,0 %	Järn (Fe)
5,0%	Mangan (Mn)
Baserat på ammonium- och kaliumsulfat, järn- och manganoxid, samt magnesiumoxid.	
Bör endast användas vid behov till golf- och fotbollsplaner, park- och trädgårdsområden. De rekommenderade mängderna får inte överskridas.	
Distributör i Sverige:	Floratine Scandinavia AB c/o Floratine Norge AS Gullfunnet 50 1570 Dilling, NORGE
Rekomenderad mängd: 15–30 g/m² (150–300 kg/ha)	
Gödsla när gräset är i tillväxt. Vattna 4-5 minuter efter påföring, för att lösa upp gödselkornen. Använd mindre gödselmängder vid höga temperaturer (> 25 °C), och vattna omedelbart för att undvika brännskador. Denna produkt innehåller järn som kan misfärga betong och andra ytor. Ta bort partiklar snabbt genom sopning eller blåsning. TVÄTTA INTE BORT MED VATTEN. I samband med gödsling bör endast torra och rena gödselspridare användas. Låt inte gödselmedel bli liggande kvar i spridaren. Gödslet skal lagras tørt.	

Norge	NK-gjødsel med Mg og Mn 8 – 0 – 13 (8 – 0 – 16)
8,0%	Nitrogen (N)
	8,0% ammonium-N
13,3%	Kalium (K) - vannløselig
	16,0% kaliumoksyd (K ₂ O)
3,0%	Magnesium (Mg)
14,5%	Svovel (S) - vannløselig
2,0%	Jern (Fe) - vannløselig
5,0%	Mangan (Mn) - vannløselig
Basert på ammonium- og kaliumsulfat, jern- og manganoksyd, samt magnesiumoksyd.	
Skal bare benyttes ved påvist behov til golf- og fotballbaner, park- og hagearealer. De anbefalte mengder må ikke overskrides.	
Distributør i Norge:	Floratine Norge AS Gullfunnet 50 1570 Dilling, NORGE Reg. nr. (Mattilsynet): 3253 Nettovekt: 20 kg
Anbefalte gjødselmengder: 10 – 30 g/m² (100 – 300 kg/ha)	
En lett oppløselig NK-gjødsel som er spesielt godt egnet til arealer som har stor fosfor-reserve i jorda, samt til arealer som har ekstra kalium-behov. Gjødsla tilføres når graset er i vekst, og på tørt bladverk. Det vannes 5-6 minutter etter at gjødsling er utført, for å løse opp gjødselkornene. Bruk mindre gjødselmengder ved høye temperaturer (> 25oC), og vann øyeblikkelig for å unngå sviskader. Dette produktet inneholder jern som kan misfarge betong og andre overflater. Fjern partikler omgående ved feing eller blåsning. VASK IKKE BORT MED VANN. I forbindelse med gjødsling skal det kun benyttes tørre og rene gjødselspredere. La ikke gjødsel bli liggende igjen i spredere. Gjødsla skal lagres tørt.	

Taiwan (R.O.C.)	8 + 0 + 16 + Mg, Fe, Mn
8.0%	Total Nitrogen (N)
	8.0% Ammoniacal Nitrogen
16.0%	Soluble Potash (K ₂ O) from potassium sulfate
3.0%	Magnesium (Mg) from magnesium oxide
14.5%	Total Sulfur (S) from sulfate
2.0%	Iron (Fe) from iron oxide
5.0%	Manganese (Mn) from manganese oxide
Application rates: 15–30 g/m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 15-30 g/m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Australia	8 + 0 + 13.3 + Mg, Fe, Mn
8.0%	Total Nitrogen (N)
	8.0% Ammoniacal Nitrogen
13.3%	Potassium (K) from potassium sulfate
1.8%	Magnesium (Mg) from magnesium oxide
14.5%	Sulfur (S) from sulfate
2.0%	Iron (Fe) from iron oxide
5.0%	Manganese (Mn) from manganese oxide
Application rates: 1.5–3.0 kg/100 m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 1.5–3.0 kg/100 m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

New Zealand	8 + 0 + 13.3 + Mg, Fe, Mn
8.0%	Total Nitrogen (N)
	8.0% Ammoniacal Nitrogen
13.3%	Potassium (K) from potassium sulfate
1.8%	Magnesium (Mg) from magnesium oxide
14.5%	Sulfur (S) from sulfate
2.0%	Iron (Fe) from iron oxide
5.0%	Manganese (Mn) from manganese oxide
Application rates: 15–30 g/m² (150–300 kg/ha)	
Directions for use: Apply product to actively growing turf at 15–30 g/m ² . Apply to dry foliage. Water in after application with 5 to 6 mm of irrigation. If temperatures exceed 25°C or if application to damp or wet turf is unavoidable, irrigate immediately after application. Apply at low rates if temperatures exceed 25°C. This product contains iron which may stain concrete and other surfaces. Remove particles promptly by sweeping or blowing to avoid staining. DO NOT WASH OFF WITH WATER. Store in a clean, dry place. Reseal opened bag by folding top down and securing. Use only clean, dry spreaders. Do not leave unused product in spreaders.	

Spreaders	Ground Speed	Width of Coverage	Spreader Settings			
			15 g/m ²	20 g/m ²	25 g/m ²	30 g/m ²
Model 2000 (cone 9); 2000 SR (cone 9) Settings for single pass application = FULL RATE	5 km/hr	3.0 m	1 3/4	J 3/4	L	M
SS-2 Drop Spreader Settings for single pass application = FULL RATE	7 km/hr	0.9 m	3 1/2	4 1/4	5	5 3/4
Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.						



Manufactured by:
The Andersons, Inc.
P. O. Box 119
Maumee, OH 43537
U.S.A.

www.andersonsplantnutrient.com

The Andersons logo and Tee Time are registered trademark of The Andersons, Inc.

20 kg/net

10009076

A24