

2,4-D, MECOPROP-P, DICAMBA	GROUP	4	HERBICIDE
SULFENTRAZONE	GROUP	14	HERBICIDE



PROFESSIONAL TURF PRODUCTS

FERTILIZER WITH SURGE®

BROADLEAF HERBICIDE



Dandelion



Chickweed

16-0-9 GRANULAR WEED AND FEED PROVIDES PLANT NUTRITION



4 active ingredients enhance speed of herbicide activity



Superior cool season performance



Controls tough broadleaf weeds

Not for sale, distribution or use in Nassau or Suffolk
Counties in New York State.

EPA Reg. No. 2217-882-9198
EPA Est. No. 9198-OH-M, 9198-OH-ZB, 9198-AL-001A
Underlined letter is first letter used in run code on bag.

ACTIVE INGREDIENTS:	
2,4-D, 2-ethylhexyl ester.....	1.153%
Mecoprop-p acid.....	0.174%
Dicamba acid.....	0.073%
Sulfentrazone.....	0.054%
OTHER INGREDIENTS.....	98.566%
TOTAL.....	100.000%

THIS PRODUCT CONTAINS:
15.30 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or
0.765% (CAS 1928-49-4)
3.48 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid
equivalent per ton or 0.174% (CAS 16484-77-8)
1.46 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.073%
(CAS 1918-00-9)
0.68 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihy-
dro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl] phenyl]
methanesulfonamide per ton or 0.034% (CAS 122836-35-5)
Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE BACK PANEL FOR COMPLETE PRECAUTIONARY STATEMENTS,
INCLUDING FIRST AID, AND DIRECTIONS FOR USE,
INCLUDING STORAGE AND DISPOSAL.

NET WEIGHT 40.00 LBS (18.14 KG)
COVERS UP TO 20,000 SQ. FT.



PROFESSIONAL TURF PRODUCTS

FERTILIZER WITH SURGE®

BROADLEAF HERBICIDE

This product is a Surge Broadleaf Herbicide, containing fertilizer, 2,4-D, mecoprop-p, dicamba and sulfentrazone. This granular weed and feed provides plant nutrients and controls a broad spectrum of listed broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.	
ACTIVE INGREDIENTS:	
2,4-D, 2-ethylhexyl ester.....	13.53%
Mecoprop-p acid.....	0.78%
Dicamba acid.....	0.073%
Sulfentrazone.....	0.034%
OTHER INGREDIENTS:	88.566%
TOTAL	100.000%
THIS PRODUCT CONTAINS:	
15.30 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.765% (CAS 1928-43-4)	
3.48 lb (+) (-)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.174% (CAS 6484-77-8)	
1.46 lb. 3,5-dichloro-p-aminic acid equivalent per ton or 0.073% (CAS 998-90-9)	
0.68 lb N-(2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-5-methyl-5-oxo-2H-2,4-hisoxal-1-yl] phenyl) methanesulfonamide per ton or 0.034% (CAS 122836-35-5)	
Isomer Specific by AOC Methods	
EPA Reg. No. 2217-882-9198	
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A	
Underlined letter is first letter used in run code on bag	

GUARANTEED ANALYSIS	
Total Nitrogen (N)	16.00%
16.00% Urea Nitrogen*	9.00%
Soluble Potash (K2O)	
Derived from urea, potassium chloride and polymer coated sulfur coated urea.	
Chlorine (Cl) Max	7.60%
*4.00% Slowly available nitrogen from polymer coated sulfur coated urea.	
F002091	

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapco.org/metals.html>

We recommend that you follow the Florida Green Industries BMP's at:
<http://www.flrules.org/Gateway/reference.asp?No=ref-04706>

We recommend that you follow the Florida Golf Course BMP's at:
<http://www.flrules.org/Gateway/reference.asp?No=ref-04702>

KEEP OUT OF REACH OF CHILDREN
CAUTION
STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
All loaders, applicators, and other handlers must wear:
• long sleeved shirt and long pants • shoes plus socks • waterproof gloves

User Safety Requirements
Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations
• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
• Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
• Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-889-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants, and to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistem or well may result in contamination of drinking water or groundwater.

Non-target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers.
- Do not apply this product to newly seeded grasses until well-established or after the third mowing.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area.
- Do not use this product on 'Floratam' St. Augustinegrass.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

WEED RESISTANCE MANAGEMENT

For resistance management, this product contains Group 4 and Group 14 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 or Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 or Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout area prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-880-5556.

WHERE TO USE

This product is a Surge® Broadleaf Herbicide containing fertilizer, 2,4-D, mecoprop-p, dicamba and sulfentrazone. This granular weed and feed provides plant nutrients and controls a broad spectrum of listed broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

This product is intended to be applied by noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations, for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrasses established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

This product may be applied to newly seeded cool season turfgrass. Apply only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the listed rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area. Do not apply product to bare ground. Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers.

California: Make broadcast applications only between March 1 and September 1. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone containing products.

New York: Only one application per year of this product is allowed. This product is not allowed to be sold or used in Nassau or Suffolk Counties.

HOW MUCH TO APPLY

For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass: Apply 150-174 pounds of this product per acre of established turfgrass. The equivalent application rates are 3.0 - 4.0 pounds of product per 1,000 square feet of turfgrass.

For New York State only: Apply 130-166 pounds of this product per acre of established turfgrass. The equivalent application rates are 3.0 - 3.8 pounds of product per 1,000 square feet of turfgrass.

For St. Augustinegrass, centipedegrass, hybrid bermudagrass, bentgrass, and colonial bentgrass: Apply 87-109 pounds of this product per acre of established turfgrass. The equivalent application rates are 2.0 - 2.5 pounds of product per 1,000 square feet of turfgrass.

Limitations on broadcast applications to ornamental turfgrass.				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Annual Rate
Ornamental Turfgrass	174 pounds/A (1.33 lb 2,4-D ae/A) (0.30 lb MCPP-p ae/A) (0.13 lb dicamba ae/A) (0.06 lb sulfentrazone/A)	2	30 Days	348 pounds/A (2.66 lb 2,4-D ae/A) (0.60 lb MCPP-p ae/A) (0.25 lb dicamba ae/A) (0.12 lb sulfentrazone/A)
ae = Acid Equivalent. Do not exceed the maximum annual rate of 348 pounds (2.66 lb 2,4-D ae, 0.60 lb MCPP-p ae, 0.25 lb dicamba ae, and 0.12 lb sulfentrazone/A) per acre per season, excluding spot treatments.				

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

APPLICATION DIRECTIONS

- Spring and fall applications are suitable for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
- Mow the turfgrass 1 to 2 days before application.
- Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
- Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
- Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

SPREADER SETTINGS			
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.			
40 lbs. treats 16,000 sq. ft. at the 109 lbs/acre - LOW RATE (2.5 lbs/1,000 sq. ft.)			
40 lbs. treats 10,500 sq. ft. at the 166 lbs/acre - MEDIUM RATE (3.8 lbs/1,000 sq. ft.)			
40 lbs. treats 10,000 sq. ft. at the 174 lbs/acre - HIGH RATE (4.0 lbs/1,000 sq. ft.)			
Spreader	Low Rate 2.5 lbs/1,000 sq. ft.	Medium Rate 3.8 lbs/1,000 sq. ft.	High Rate 4.0 lbs/1,000 sq. ft.
The Andersons LCO-1000	K	L 3/4	M
The Andersons Yard Star 2150	13	14 1/2	15
Agri-Fab Broadcast (1-10)	3 1/4	5	5 1/4
Brinly 20 Series Push Broadcast (0-30)	13	14 1/2	16
Chapin Broadcast	13	14 1/2	16
EarthWay Broadcast (0-30)	13	14 1/2	16
Echo Broadcast	3 3/4	5 1/2	5 3/4
Lesco Broadcast	10	12	12 1/2
Scotts Broadcast (2-15)*	3 3/4	5 1/2	5 3/4

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

DISTRIBUTION AND SPREADER PATTERNS

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

WEEDS CONTROLLED

Dandelion, chickweed, plantain, and clover are controlled effectively with this product.

A list of susceptible weeds is shown below.

- BROADLEAF WEEDS -			
Annual yellow sweetclover	Carpetweed	Hairy fleabane	Poison oak
Aster	Catchweed bedstraw	Hawkweed	Poorjoe
Austrian fieldress	Catnip	Healall	Prairie sunflower
Bedstraw	Catsear	Heartleaf drymaria	Prostrate knotweed
Beggaricks	Chickweed, common	Henbit	Prostrate pigweed
Beltony, Florida	Chickweed, mouseear	Hoary cress	Prostrate spurge
Bird vetch	Chickweed, mouseear	Hoary plantain	Puncturevine
Bitter wintercress	Chickweed, mouseear	Hoary vervain	Purslane, common
Bitterness, hairy	Chicory	Knawel	Ragweed
Bitterweed	Cinquefoil	Kniveyed	Red sorrel
Black-eyed Susan	Clover, crimson	Lambsquarters	Redroot pigweed
Black medic	Clover, hop	Lespedeza	Redstem filaree
Blackseed plantain	Clover, red	Mallow	Rough cinquefoil
Blessed thistle	Clover, strawberry	Matchweed	Russian pigweed
Bloodflower	Clover, sweet	Mexicanweed	Scarlet pimpernel
Blue milkweed	Clover, white	Milk vetch	Sheep sorrel
Blue lettuce	Cockle	Mouseear	Sheepspurge
Blue vervain	Creeping jenny	hawkweed	Slender plantain
Bracted plantain	Cudweed	Mugwort	Smallflower
Bristly oxtongue	Curly dock	Narrowleaf plantain	gallinsoya
Broadleaf dock	Daisy, English	plantain	Smooth dock
Broadleaf plantain	Daisy fleabane	Nettle	Smooth pigweed
Buckhorn plantain	Daisy, oxeve	Orange hawkweed	Sorrel
Bulbous buttercup	Dandelion	Onails	Sowthistle
Bull thistle	Dichondra	Parsley-piert	Spanishneedles
Bullnettle	Dollarweed	Parsnip	Speedwell
Burclover	False dandelion	Pearlwort	Spiny amaranth
Burdock	Falsellax	Pepperweed	Spiny sowthistle
Burning nettle	False sunflower	Pigweed	Spotted catsear
Burweed	Florida pusley	Pineywoods	Spotted spurge
Buttercup	Gallinsoga	bedstraw	Sourweed
Buttonweed	Goldrood	Plains coreopsis	Strawberry, India
Canada thistle	Ground ivy	Plantain	mock
Carolina geranium	Gumweed	Poison ivy	Tall nettle

STORAGE & DISPOSAL	
Do not contaminate water, food, or feed by storage and disposal.	
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.	
PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representatives at the nearest EPA Regional Office for guidance.	
CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.	

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Distributed By:
The Andersons, Inc.
PO Box 119 | Maumee, OH 43537

©2014 The Andersons, Inc. All rights reserved. The Andersons logo is a registered trademark of The Andersons, Inc. Surge is a registered trademark of F&B/Gordon Corporation.

10008873
30001810
F24

10008873
30001810
F24
The Andersons Professional is up to 20,000 sq. ft. (40 lb)