



THE ANDERSONS PROFESSIONAL TURF PRODUCTS

FERTILIZER 16-0-8 WITH ESCALADE[®] HERBICIDE

- This product is a premium grade herbicide and turf food combination which controls a wide spectrum of broadleaf weeds while providing both fast and long feeding action.
- Contains the esters of 2,4-D and Fluroxypyr with Dicamba acid.
- Weeds contacted by this product gradually die and disappear from the lawn.

ACTIVE INGREDIENTS:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	0.8650%
1-Methylheptyl Ester of Fluroxypyr: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic Acid, 1-methylheptyl Ester**	0.2060%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.0717%
OTHER INGREDIENTS:	98.8573%
Total	100.0000%

Isomer Specific Method, Equivalent to:

* 2,4-Dichlorophenoxyacetic Acid	0.5740%
** ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic Acid	0.1430%
***3,6-Dichloro-o-Anisic Acid	0.0717%

GUARANTEED ANALYSIS

Total Nitrogen (N)	16.00%
16.00% Urea Nitrogen*	
Soluble Potash (K ₂ O)	8.00%
Derived from urea, potassium chloride and polymer coated sulfur coated urea.	
Chlorine (Cl) Max	6.78%

*4.00% Slowly available nitrogen from polymer coated sulfur coated urea.

Information regarding the contents and levels of metals in this product is available on the internet at <https://www.aapfco.org/metals.html>

EPA Reg. No. 228-427-9198

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code.

Do not sell, distribute or use this product in Nassau and Suffolk counties, New York

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

Net Wt. 40 Lbs. (18.14 kg)

THE ANDERSONS PROFESSIONAL TURF PRODUCTS

FERTILIZER 16-0-8 WITH ESCALADE® HERBICIDE

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves made of any waterproof material, selection category A.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Alder	Broadleaf plantain	Coffeeweed	Field pansy	Hop clover
Annual yellow	Broomweed	Common	Flea bane (daisy)	Horse-nettle
sweet clover	Buckhorn	chickweed	Floweed	Horse-tail
Artichoke	Buckhorn plantain	Common mullein	Florida betony	Indiana mallow
Aster	Bulbous buttercup	Common	Florida pusley	Ironweed
Austrian fieldcress	Bull nettle	sowthistle	Frenchweed	Jewelweed
Bedstraw	Bull thistle	Corn Chamomile	Galinsoga	Jimsonweed
Beggar-tick	Burdock	Creeping jenny	Garlic mustard	Kochia
Biden	Burning nettle	Crimson clover	Goatshead	Knawel
Bindweed	Bur ragweed	Croton	Goatsbeard	Knotweed
Bird vetch	Burweed	Cudweed	Goldenrod	Lambsquarter
Bitterweed	Buttercup	Curly dock	Ground ivy	Lespedeza
Bitter wintercress	Canada thistle	Curly indigo	Gumweed	Locoweed
Black-eyed Susan	Carolina geranium	Dandelion	Hairy bittercress	Lupine
Black medic	Carpetweed	Dead nettle	Hairy fleabane	Mallow
Black mustard	Catchweed	Dock	Hawkweed	Marshelder
Black-seed	bedstraw	Dogbane	Healall	Matchweed
plantain	Catsear	Dogfennel	Heartleaf drymary	Mexicanweed
Blessed thistle	Catnip	Elderberry	Hedge bindweed	Milk vetch
Blue lettuce	Chickweed	English daisy	Hedge mustard	Milkweed blood
Blue vervain	Chicory	Fall dandelion	Hemp	flower
Box elder	Cinquefoil	False dandelion	Henbit	Mugwort
Bracted plantain	Clover	False flax	Hoary cress	Morningglory
Brassbuttons	Cockle	False sunflower	Hoary plantain	Mouseear
Bristly oxtongue	Cocklebur	Fiddleneck	Hoary vervain	chickweed
Broadleaf dock	Coffeebean	Field bindweed	Honeysuckle	Musk thistle

Mustard	Primrose	Smooth dock	Tarweed	Wild four-o'clock
Narrowleaf	Prostrate	Smooth pigweed	Thistle	Wild garlic
plantain	knottweed	Sneezeweed	Tick trefoil	Wild geranium
Narrowleaf vetch	Prostrate pigweed	Southern wild	Toadflax	Wild lettuce
Nettle	Prostrate spurge	rose	Trailing	Wild marigold
Orange	Prostrate vervain	Sowthistle	Crownvetch	Wild onion
hawkweed	Puncture vine	Spanishneedle	Tumble mustard	Wild parsnip
Oxalis	Purslane	Spatterdock	Tumble pigweed	Wild radish
Oxeye daisy	Ragweed	Speedwell	Tumbleweed	Wild rape
Parsley-piert	Red clover	Spiny Amaranth	Velvetleaf	Wild strawberry
Parsnip	Redroot pigweed	Spiny cocklebur	Venice mallow	Wild sweet potato
Pearwort	Red sorrel	Spotted catsear	Veronica	Wild vetch
Pennycress	Redstem filaree	Spotted	Vervain	Willow
Pennywort	Rough cinquefoil	knawweed	Vetch	Witchweed
Peppergrass	Rough fleabane	Spotted spurge	Virginia	Woolly
Pepperweed	Roundleafed	Spurge	buttonweed	morningglory
Pineywoods	marigold	Spurweed	Virginia creeper	Woodsorrel
bedstraw	Rush	Stinging nettle	Virginia	Woolly croton
Plains coreopsis	Russian pigweed	Stinkweed	pepperweed	Woolly plantain
Plantain	Russian thistle	Stitchwort	Wayleaf	Wormseed
Poison hemlock	St. Johnswort	Strawberry dower	bullthistle	Yarrow
Poison ivy	Scarlet pimpernel	Sumac	Western clematis	Yellow rocket
Poison oak	Scotch thistle	Sunflower	Western salsify	Yellowflower
Pokeweed	Sheep sorrel	Sweet clover	White clover	Pepperweed
Poorjoe	Sheepspurse	Tall nettle	White mustard	and other
Povertyweed	Slender plantain	Tall vervain	Wild mustard	broadleaf
Prickly lettuce	Smallflower	Tansy mustard	Wild aster	weeds
Prickly sida	galinsoga	Tansy ragwort	Wild buckwheat	
	Smartweed	Tanweed	Wild carrot	

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 40 pound bag should be used on 10,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are **approximate** for applying contents of this package at a suggested rate of 4 lbs. per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 lb. per 250 square feet.

SPREADER SETTINGS			
One pass application			
Suggested spreader settings are not intended to replace calibration.			
Please calibrate your spreader prior to applying product.			
40 lbs. treats 10,000 sq. ft. at the 174 lbs/acre rate.			
Spreader	Ground Speed	Width of Coverage	Spreader Settings 4.0 lbs/1,000 sq. ft.
Andersons 2000 Series (cone G)	3 mph	8 feet	K
LESCO 020093 (0.25)	3 mph	6 feet	E 1/4
LESCO 705698 (0.25)	3 mph	6 feet	12 1/2
Scotts R-8A (cone G)	3 mph	8 feet	1 1/2
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm)	4.5 mph	26 feet	4 3/4 II-C
Vicon (O3 series)	4.5 mph	26 feet	25

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound of nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *0.64 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

Limitations: The maximum rate per application is 6 pounds 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 12 pounds per 1,000 square feet, excluding spot treatments.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material.

Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, or offered to others for their use, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, BUYERS OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any directions for use, the warranty disclaimer, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. (RWD02610)

Distributed by:
The Andersons, Inc.
P.O. Box 119
Maumee, OH 43537
www.AndersonsPro.com

The Andersons®

© 2025 The Andersons, Inc. All rights reserved. The Andersons logo is a registered trademark of The Andersons, Inc. Escalade is a registered trademark of Nufarm America, Inc. R25

10005603
30001819

